

Monoclonal Mouse Antibody to Hepatocyte Specific Antigen

Catalog No.:	PDM 162
Intended Use:	This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Clinical interpretation of staining results should be accompanied by histological studies with proper controls. Patients' clinical histories and other relevant diagnostic tests should be utilized by a qualified person(s) when evaluating and interpreting results.
Immunogen:	Formalin-fixed, failed allograft human liver
Clone:	OCH1E5 (also known as Hep Par 1)
Isotype:	IgG1, kappa
Format:	This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded and acetone fixed cryostat tissue sections. No further titration is required.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. <u>Formalin fixed tissues do not require any special treatment; however, heat-induced epitope retrieval with 10 mM Citrate, pH 6.0 or 1 mM EDTA, pH 8.0 may increase signal detected.</u>
Specificity:	Hepatoblastoma is the most common primary tumor of the liver in children. Use of specific hepatocyte markers and alpha-fetoprotein or carcinoembryonic antigen are useful for identifying normal and malignant fetal hepatocytes. This antibody recognizes an uncharacterized antigen present in both adults and fetal normal hepatocytes to produce distinct granular cytoplasmic staining. This antibody stains the majority of hepatocellular carcinomas.
Positive Control:	Liver
Cellular Localization:	Cytoplasmic
Storage:	Store at 2-8°C. Do not use beyond the expiration date stated on the label.
References:	(i) Amarapurkar et al. Indian J Pathol Microbiol 49 (3): 341, 2006. (ii) Chu et al. Am J Surg Pathol 26 (8): 978, 2002. (iii) Fasano et al. Mod Pathol 11: 934, 1998. (iv) Lamps and Folpe Adv Anat Pathol 10 (1): 39, 2003. (v) Siddiqui et al. Cancer Cytopathol 96 (1): 49, 2002. (vi) Wennerberg et al. Am J Pathol 143: 1050, 1993.

IVD: For In Vitro Diagnostic Use

DBS will not be held responsible for patent infringement or other violation that may occur with the use of our product

DBS

1020 Serpentine Lane, # 114, Pleasanton, CA 94566 Tel: 925 484 3350, Fax: 925 484 3390

Website: www.dbiosys.com e-mail: customersupport@dbiosys.com