

# **IHC Made Affordable**

# Montage<sup>™</sup> Tween 20

Catalog Number: K067-125AN

Document #:	DS-2041-B
Effective Date:	12/18/2023

# Intended Use

For In Vitro Diagnostic Use.

# **Product Description**

Tween 20 is a detergent (surfactant) commonly added to buffers and reagents for immunohistochemistry. Its purpose is to decrease background staining and enhance reagent spreading in automated and manual procedures.

# **Summary and Explanation**

Tween-20 is a non-ionic surfactant that is useful for solubilizing membrane proteins during isolation. Tween-20 is also added to buffers for immunohistochemistry. Tween-20 may also help to prevent non-specific antibody binding and function as a blocking agent in immunoblotting assays.

# Format

Concentrated

#### Volume/UOM

125 mL

# **Storage and Handling**

Store at room temperature. Do not use after expiration date printed on label.

# **Preparation of Working Solutions**

1. It is suggested to use the product at concentrations between 0.05% and 0.1% in buffers and reagents.

# **Protocol Recommendations**

1. Tween 20 is a detergent (surfactant) commonly added to buffers and reagents for immunohistochemistry.

# **Quality Control**

Refer to CLSI Quality Standards for Design and Implementation of Immunohistochemistry Assays; Approved Guideline-Second edition (I/LA28-A2). CLSI Wayne, PA, USA (www.clsi.org). 2011.

# Troubleshooting

Contact Diagnostic BioSystems Technical Support at (925) 484-3350, extension 2, techsupport@dbiosys.com or your local distributor to report unusual staining results.

# Warranty

There are no warranties, expressed or implied, which extend beyond this description. Diagnostic BioSystems is not liable for property damage, personal injury, or economic loss caused by this product.

# **Performance Characteristics**

The protocols for a specific application can vary. These include, but are not limited to: fixation, heat-retrieval method, incubation times, tissue section thickness and detection kit used. Due to the superior sensitivity of these unique reagents, the recommended incubation times and titers listed are



6616 Owens Drive Pleasanton, CA 94588 Tel: (925) 484 3350 www. dbiosys.com

**Diagnostic BioSystems** 





Gotthardstrasse 28 6302 Zug Switzerland



MedEnvoy Global B.V. Prinses Margrietplantsoen 33 - Suite 123 2595 AM The Hague The Netherlands

not applicable to other detection systems, as results may vary. The data sheet recommendations and protocols are based on exclusive use of Diagnostic BioSystems products. Ultimately, it is the responsibility of the investigator to determine optimal conditions. These products are tools that can be used for interpretation of morphological findings in conjunction with other diagnostic tests and pertinent clinical data by a qualified pathologist.

# Precautions

This product is a single-use, non-sterile, in vitro diagnostic device.

- 1. Wear disposable gloves when handling reagents.
- 2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come in contact with sensitive areas, wash with copious amounts of water.
- 3. Microbial contamination of reagents may result in an increase in nonspecific staining.
- 4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.
- 5. Do not use reagent after the expiration date printed on the label.
- 6. The MSDS is available upon request.
- Consult OSHA, federal, state or local regulations for disposal of any toxic substances.